

# **Handbook of Small Animal Regional Anesthesia and Analgesia Techniques**

# Handbook of Small Animal Regional Anesthesia and Analgesia Techniques

## **Phillip Lerche BVSc PhD DipACVAA**

Associate Professor – Clinical  
Anesthesiology and Pain Management  
Department of Veterinary Clinical Sciences  
The Ohio State University  
College of Veterinary Medicine  
Columbus, Ohio, USA

## **Turi K. Aarnes DVM MS DipACVAA**

Associate Professor – Clinical  
Anesthesiology and Pain Management  
Department of Veterinary Clinical Sciences  
The Ohio State University  
College of Veterinary Medicine  
Columbus, Ohio, USA

## **Gwen Covey-Crump BVetMed CertVA DipECVAA MRCVS**

Specialist in Veterinary Anaesthesia and Analgesia  
Clinical Teacher  
Langford Veterinary Services  
University of Bristol  
Bristol, UK

## **Fernando Martinez Taboada LV CertVA DipECVAA MRCVS**

Specialist in Veterinary Anaesthesia and Analgesia  
The Veterinary Teaching Hospital  
Faculty of Veterinary Science  
The University of Sydney  
Sydney, Australia

**WILEY** Blackwell

This edition first published 2016 © 2016 by John Wiley & Sons, Ltd

*Registered office:* John Wiley & Sons, Ltd, The Atrium, Southern Gate, Chichester,  
West Sussex, PO19 8SQ, UK

*Editorial offices:* 9600 Garsington Road, Oxford, OX4 2DQ, UK  
The Atrium, Southern Gate, Chichester, West Sussex, PO19 8SQ, UK  
1606 Golden Aspen Drive, Suites 103 and 104, Ames, Iowa 50010, USA

For details of our global editorial offices, for customer services and for information about how to apply for permission to reuse the copyright material in this book please see our website at [www.wiley.com/wiley-blackwell](http://www.wiley.com/wiley-blackwell)

The right of the author to be identified as the author of this work has been asserted in accordance with the UK Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the publisher.

Designations used by companies to distinguish their products are often claimed as trademarks. All brand names and product names used in this book are trade names, service marks, trademarks or registered trademarks of their respective owners. The publisher is not associated with any product or vendor mentioned in this book. It is sold on the understanding that the publisher is not engaged in rendering professional services. If professional advice or other expert assistance is required, the services of a competent professional should be sought.

The contents of this work are intended to further general scientific research, understanding, and discussion only and are not intended and should not be relied upon as recommending or promoting a specific method, diagnosis, or treatment by health science practitioners for any particular patient. The publisher and the author make no representations or warranties with respect to the accuracy or completeness of the contents of this work and specifically disclaim all warranties, including without limitation any implied warranties of fitness for a particular purpose. In view of ongoing research, equipment modifications, changes in governmental regulations, and the constant flow of information relating to the use of medicines, equipment, and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each medicine, equipment, or device for, among other things, any changes in the instructions or indication of usage and for added warnings and precautions. Readers should consult with a specialist where appropriate. The fact that an organization or Website is referred to in this work as a citation and/or a potential source of further information does not mean that the author or the publisher endorses the information the organization or Website may provide or recommendations it may make. Further, readers should be aware that Internet Websites listed in this work may have changed or disappeared between when this work was written and when it is read. No warranty may be created or extended by any promotional statements for this work. Neither the publisher nor the author shall be liable for any damages arising herefrom.

A catalogue record for this book is available from the Library of Congress and the British Library.

ISBN: 9781118741825

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic books.

Main cover image: with permission from The Ohio State University

Set in 9/12pt Meridien LT Std by Aptara Inc., New Delhi, India

# Contents

## **1 Introduction, 1**

- Reasons to provide regional anesthesia, 2
  - History of regional anesthesia/analgesia, 3
  - Principles of the major techniques, 3
  - Local anesthetic drugs, 4
  - Additives to local anesthetics, 6
- Equipment, 7
  - Syringes, 7
  - Needles, 7
  - Epidural catheter kits, 8
  - Nerve locators, 8
  - Differential pressure transducer, 9
  - Acoustic device, 9
- Complications and contraindications, 9
  - Local, 10
  - Systemic, 10
- Species differences and considerations, 10
- Safe practice, 11
  - Gaining confidence, 11
  - Making a plan, 11
- References, 11
- Further reading, 12

## **2 Cutaneous Innervation Index, 13**

- Cutaneous innervation index, 14
  - Canine dermatomes
    - Body, 14
    - Head, 15
    - Thoracic limb, 16
    - Pelvic limb, 18

### **3 Infiltration Blocks, 21**

Infiltration for mass removal, 22

Line and ring blocks, 22

Inverted “L” block, 24

Block of the surgical incision, 26

Single injection of the surgical incision , 26

Diffusion catheter/“soaker” catheter/“pain buster”, 28

Intercostal blocks, 30

Interpleural block (also known as intrapleural or pleural), 32

Intraperitoneal block, 34

References, 35

### **4 Blocks of the Head, 37**

Retrobulbar, 38

Maxillary, 40

Infraorbital, 42

Mandibular (or inferior alveolar), 44

Mental (or middle mental), 46

Auriculotemporal and great auricular, 48

References, 52

### **5 Regional Anesthetic Blocks of the Limbs, 53**

Thoracic limb regional anesthetic blocks, 54

Brachial plexus, 56

Paravertebral: C6–T1, 58

RUMM (radial/ulnar/median/musculocutaneous), 60

Bier block (intravenous regional anesthesia), 62

Pelvic limb, 64

Femoral/saphenous nerve block, 66

Sciatic nerve block, 70

References, 74

### **6 Epidurals and Spinals, 75**

Lumbosacral epidural and spinal in dogs, 76

Lumbosacral epidural in cats, 78

Caudal epidural in dogs and cats, 80

References, 82

### **7 Emergency Procedures, 83**

Complications associated with local anesthetic techniques, 84

Bleeding, 84

Nerve trauma, 84

Complications associated with local anesthetic drugs, 84

    Allergic reactions to local anesthetic drugs, 84

    Methemoglobinemia, 85

    Systemic toxicity of local anesthetics, 85

    Cardiovascular toxicity, 85

References, 88

Further reading, 88

Index, 89

## CHAPTER 1

# Introduction

Phillip Lerche

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.  
By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.  
© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## Reasons to provide regional anesthesia

The use of regional anesthesia as a component of perioperative pain management has gained acceptance and popularity in small animal practice over the past few decades. Reasons for this include the fact that many of the regional blocks are straightforward to perform, requiring moderate technical skill given familiarity with patient anatomy; they can be conducted relatively safely given an understanding of local anesthetic drug pharmacology, complications and side effects; and they contribute to the two major tenets of treating pain: pre-emptive and multimodal analgesia.

Providing pre-emptive analgesia by performing regional anesthesia prior to surgery leads to a drastic reduction in intraoperative nociceptive (pain) stimulation. This results in a decrease in anesthetic maintenance drug as well as intra- and postoperative analgesic requirements, thereby decreasing the incidence of drug side effects during surgery, and improving postoperative patient comfort as well as duration of pain relief. Some techniques can be continued postoperatively to assist in managing pain after particularly painful surgeries once the patient has recovered from anesthesia, e.g. instilling local anesthetic into a chest tube after thoracotomy, or injecting local anesthetic into an epidural or spinal catheter after pelvic limb or abdominal surgery.

The experience of pain, a sensory process involving the nociceptive pathway, is complex, and involves several steps. Noxious stimuli involving mechanical, chemical or thermal injury to tissue are first transduced into electrical stimuli by peripheral nociceptors (pain receptors). These electrical impulses are then transmitted to the spinal cord, where they are modulated by neurons in the dorsal horn of the gray matter of the spinal cord. Here, impulse intensity can be increased (amplified) or decreased (suppressed). Finally, the nociceptive signals are projected via lateral nerve fibers to the brain where they are perceived.

Whereas most analgesic drugs either decrease the amount of excitatory neurotransmitters, or increase the level of inhibitory neurotransmitters released in the nociceptive pathway, drugs used to provide regional anesthesia block sodium channels in neurons. This completely prevents sensory neurons from transmitting noxious stimuli from the periphery to the brain and spinal cord, or from the spinal cord to the brain in the case of epidural or spinal analgesia, thus providing effective pain relief for the duration of the block. Using regional anesthetic techniques in conjunction with other analgesic drugs that act in different ways on the nociceptive fibers (e.g. with opioids, alpha-2 agonists, ketamine) results in multimodal analgesia, contributing to an overall decrease in excitatory neurotransmission within



the pain pathway both during and after surgery. This approach allows for the lowest effective dose of each drug to be used, which decreases side effects and enhances patient safety.

## **History of regional anesthesia/analgesia**

The use of a local anesthetic drug was first demonstrated in 1884 when cocaine was used to desensitize the eye prior to surgery. Due to cocaine being habit forming and having a low safety margin, as well as the emergence of techniques allowing artificial synthesis of chemical compounds in the 1900s, non-toxic, non-addictive local anesthetics were sought, discovered, and manufactured. Initially, amino ester-type compounds were produced, until in 1943 lidocaine, an amino amide drug, was developed. Amide-type local anesthetic drugs are preferred for their longer duration of action, and several compounds in this group were discovered in the latter half of the 20th century, including mepivacaine, bupivacaine, and ropivacaine.

## **Principles of the major techniques**

### ***Topical application***

Local anesthetic drops, e.g. proparacaine, can be directly applied to the eye for immediate relief of keratoconjunctival pain, although prolonged use delays corneal healing and is not recommended. Local anesthetic can also be directly applied to exposed tissue. Drug is directly deposited into the surgical field by dripping from a syringe, or soaking surgical sponges in local anesthetic and applying them to the tissue. Local anesthetic can also be instilled into the chest via a thoracostomy tube to desensitize the pleura following chest surgery, and into the abdominal cavity to treat pain following incision of the peritoneum. Local anesthetic cream is available as a mixture of lidocaine and prilocaine, which is used to desensitize skin for intravenous catheter placement. Lidocaine is also available as a transdermal patch.

### ***Regional infiltration***

Continuous regional analgesia is accomplished by placing fenestrated “soaker” catheters in areas that are not amenable to peripheral or regional analgesic techniques. The catheter is then attached to an infusion pump or an elastomeric bulb which delivers a set rate of local anesthetic over a specified period of time.

### ***Intravenous regional analgesia***

Analgesia can be provided to a distal limb by placing an esmarch bandage and injecting local anesthetic (lidocaine only) into a vein.

### ***Intra-articular injection***

Local anesthetics injected into joints have a long duration of action due to slow systemic uptake. There is *in vitro* evidence that local anesthetics may be detrimental to chondrocyte health, with preservative-free formulations being preferred. However, *in vivo*, this has not been shown to be definitively the case (Chu et al, 2008).

### ***Peripheral nerve blockade***

Individual or groups of sensory nerves supplying a specific region are located by palpation, electrophysiology, ultrasound or varying combinations of two or three of these techniques. Local anesthetic is then deposited adjacent to, but not into, the nerves. Nerves are typically blocked at sites proximal and distant to the site of surgery.

### ***Epidural and spinal injection***

Using specific epidural/spinal needles, local anesthetic is deposited either into the extradural space (epidural injection) or into the subarachnoid space (spinal or intrathecal injection). This provides longer lasting, more intense analgesia and muscle relaxation, while minimizing systemic side effects. Preservative-free formulations of drug are recommended for epidural or spinal injection whenever possible.

## **Local anesthetic drugs**

### ***Mechanism of action***

Local anesthetics mainly act by blocking sodium ( $\text{Na}^+$ ) channels, which prevents depolarization of the neuronal cell membrane, and thus generation of an electrical impulse does not occur in response to noxious stimuli. There is evidence to suggest that local anesthetics can also exert their activity by blocking calcium channels and inhibiting reuptake of the inhibitory neurotransmitter GABA, thus enhancing its effect.

### ***Physicochemical properties***

Local anesthetic drugs have an aromatic ring and an amine group separated by a hydrocarbon chain. The amine group can be ester or amide linked. Esters are typically shorter acting than amides as they can be hydrolyzed by plasma cholinesterases. Esters are therefore not reliant on the liver for clearance, whereas amides undergo hepatic metabolism. Speed of onset is inversely proportional to the drug's lipid solubility and  $\text{pK}_a$ , i.e. the pH at which the drug exists in equal amounts of charged and non-charged molecules. Duration of effect increases as lipid solubility increases, and decreases as the rate of systemic absorption increases. Drugs that cause vasodilation, like lidocaine, have a shorter duration of action.

**Table 1.1** Clinical pharmacology of local anesthetic agents in cats and dogs

	<b>Lidocaine</b>	<b>Mepivacaine</b>	<b>Bupivacaine</b>	<b>Ropivacaine</b>
Onset (minutes)	5–10	5–10	20–30	20–30
Duration (hours)	1–3	1.5–3	3–8	3–8
Clinical dose (mg/kg)	0.5–2.0 dog 0.5–1.5 cat	Up to 3.0 dog Up to 1.5 cat	1.0–1.5 dog 1.0 cat	Up to 3.0 dog*
Toxic dose (mg/kg)	6.0 dog 3.0 cat	6.0 dog 3.0 cat	3.0 dog 2.0 cat	5.0 dog*

\*Toxic dose not established in cats. Recommend not exceeding 2 mg/kg total dose.

### Specific drugs

Amide-type local anesthetic drugs are preferred in current veterinary practice for their longer duration of action compared to ester-type drugs. See Table 1.1 for summary information.

#### Lidocaine

Lidocaine has a short onset due to its low  $pK_a$  of 7.9. Duration of action is short, lasting up to 2 h. This is due to its relatively low level of protein binding (70%), and the fact that it is a potent vasodilator. Lidocaine formulated with epinephrine has a longer duration of action due to the vasoconstriction epinephrine causes. Lidocaine is less toxic than other amide-type drugs if administered intravenously (IV), and can be administered IV to treat pain systemically, as well as to treat ventricular dysrhythmias.

#### Mepivacaine

Mepivacaine has a low  $pK_a$  of 7.6, and therefore a rapid onset of action. It is highly protein bound (95%), resulting in a duration of action of 6–8 h.

#### Bupivacaine

Bupivacaine has a  $pK_a$  of 8.1 and is highly protein bound (95%), resulting in a longer duration of action (6–8 h). The margin of safety is the lowest when compared to lidocaine, mepivacaine, and ropivacaine.

#### Ropivacaine

Ropivacaine has physicochemical properties similar to bupivacaine, and therefore has a similar onset and duration of action. Other local anesthetics are synthesized as racemic mixtures, whereas ropivacaine is a pure S-enantiomer, and has a wider margin of safety than bupivacaine.

### **Combination of drugs**

Lidocaine and bupivacaine can be mixed in a 1:1 ratio to take advantage of lidocaine's shorter onset and bupivacaine's longer duration of action.

### **Additives to local anesthetics**

Other drugs can be added to local anesthetics to enhance or extend blockade, or to decrease the pain experienced on injection.

#### ***Opioids***

Opioids, particularly preservative-free morphine, are commonly administered with local anesthetics for epidural or spinal analgesia. This produces additive or synergistic multimodal analgesic effects, with analgesia lasting up to 24 h, far longer than with individual drug therapy. Complications of adding opioids include respiratory depression, particularly if high doses or volumes are used, urinary retention, vomiting (in conscious dogs), and pruritus. Myoclonus, hindlimb paresis, altered proprioception, and hyperesthesia are rare complications.

#### ***Alpha-2 adrenoceptor agonists***

Agonism of pre- and postsynaptic alpha-2 receptors in the pain pathway results in analgesia. Xylazine (0.25 mg/kg) and medetomidine (15 µg/kg) have been used epidurally in dogs, with medetomidine providing analgesia for up to 8 h. Side effects included bradycardia and hypertension. Dexmedetomidine produces analgesia in a dose-dependent manner when given intrathecally and epidurally in dogs.

#### ***Ketamine***

Ketamine most likely produces its analgesic effects when administered epidurally by blocking NMDA channels. Ketamine also blocks some sodium and potassium channels, thus decreasing propagation of nociceptive signals. Effective doses are 1–3 mg/kg. Side effects include increased heart rate, blood pressure, and myocardial work.

#### ***Epinephrine (adrenaline)***

Epinephrine can be added to local anesthetic at a concentration of 5 µg/mL. Deposition of the combination results in local vasoconstriction, which leads to decreased systemic uptake and prolongation of blockade. This delayed uptake also results in fewer systemic side effects due to decreased plasma concentration of local anesthetic. Epinephrine should not be used when performing a Bier block or a ring block, as nerve ischemia may occur.

### ***Sodium bicarbonate***

Addition of sodium bicarbonate at 1 mEq per 10 mL of local anesthetic increases the amount of active (non-ionized) drug present, increasing diffusion across the cell membranes of neurons. This may lead to a shorter onset and longer duration of blockade. Pain experienced on injection of local anesthetic in the conscious patient is also decreased by addition of sodium bicarbonate. Sodium bicarbonate should not be added to bupivacaine or ropivacaine as precipitation will occur.

### ***Hyaluronidase***

Hyaluronidase improves permeability of tissue by depolymerizing hyaluronic acid, resulting in better spread of local anesthetic. It can be added at 3.75 IU per mL of local anesthetic to enhance the quality of blockade. Due to the enhanced permeability, duration may be decreased and toxicity increased due to increased systemic absorption. Adding hyaluronidase to ropivacaine does not enhance spread.

## **Equipment**

### **Syringes**

Due to the variety of sizes of dogs, a variety of sizes of syringes ranging from 1 mL to 20 mL should be available. Glass syringes, or specially made low-resistance plastic syringes, are useful when performing epidural techniques to check for loss of resistance. A specific type of syringe, called an Episure<sup>TM</sup> syringe, has been evaluated in people as an alternative to glass syringes when performing epidural anesthesia. This syringe features a compression spring, which supplies a constant pressure when attached to a needle. The operator can advance the needle using both hands, rather than one hand, allowing for steadier needle advancement and a visual sign that the epidural space has been entered (Riley & Carvalho, 2007).

### **Needles**

Hypodermic needles ranging from 25 Ga to 20 Ga and 2–5 cm are used to perform many local nerve blocks.

Epidural/spinal needles are used when performing epidural or spinal analgesia. These needles have a sharp bevel and incorporate a stylet to avoid depositing tissue cores within the epidural or spinal space. These needles are available in a range of gauges (18, 20, and 22 Ga) and lengths (3.8, 6.3, and 9.0 cm).

A modified spinal needle, or Tuohy needle, is slightly curved at the tip, which facilitates advancement of the catheter when placing it in

the epidural space. This type of needle is usually marked in increments of 1 cm along its length to assist with determining how far to insert the catheter. Tuohy needles are typically of larger gauge (16 and 18 Ga) than standard epidural needles in order to place the catheter more easily, although this may hinder placement in small patients.

### **Epidural catheter kits**

Sterilized kits for placing epidural catheters are available commercially. They typically include a Tuohy needle, a loss of resistance syringe, a radiopaque catheter with guidewire, a connector, and an antibacterial filter. Some kits also include a sterile pen, which is used to mark on the catheter the distance it should be advanced through the Tuohy needle.


### **Nerve locators**

Nerve locators are used to improve accuracy of local anesthetic deposition, decrease local anesthetic dose required, and limit side effects.

### **Peripheral nerve stimulators**

Peripheral nerve stimulators (PNS) are used in conjunction with Teflon-coated needles, which have a small conductive area at the tip, to locate nerves using an electrical current (Figure 1.1). The Teflon coating ensures that only the tip of the needle transmits the electrical current so that a high current density is achieved. The reader is advised to refer to the instruction leaflet enclosed with the peripheral nerve stimulator.

Briefly, the positive lead (anode) is attached to the skin by means of a sticky electrode (ECG pad) and the negative lead (cathode) is connected to the needle. Anatomical landmarks are palpated and the needle is inserted through the skin. The needle is advanced until it is in close proximity to the nerve. The operator then turns the nerve stimulator on, setting a current of 1.0–2.0 mA, pulse frequency 1–2 Hz and



**Figure 1.1** A, Peripheral nerve stimulator. B, Teflon-coated needle. C, Positive lead (anode) and sticky electrode (ECG pad). D, Negative lead (cathode). E, Syringe, injection port, and tubing. Reproduced with permission from The Ohio State University.

duration 0.1–0.3 msec. This generates an electrical field adjacent to the nerve, resulting in depolarization and muscle contraction, or twitches. The current required to elicit a twitch increases exponentially as the distance between the nerve and the needle increases. The current is reduced to the lowest possible setting required to elicit a twitch, and this is done whilst carefully redirecting the needle so that it is in close proximity to the nerve. The useful current is not a painful stimulus, and can be used in a sedated patient. Once the lowest possible current required to elicit a twitch has been determined, local anesthetic is injected through the needle via tubing. Injection of local anesthetic pushes the needle a little further away from the nerve, so twitches are usually diminished or lost immediately after injection.

### **Ultrasound**

Ultrasound probes can be used to visualize nerves directly both prior to and during needle placement and anesthetic drug deposition, which allows for improved accuracy of blockade. This technique is often combined with a nerve stimulator to confirm nerve location, allowing drug deposition close to, but not into, the nerve.

### **Differential pressure transducer**

The likelihood of injury to nerves during local anesthetic injection is increased when injections are performed under high pressure. The use of an in-line pressure transducer allows consistent force to be applied during injection.

### **Acoustic device**

Identification of needle entry into, or puncture of, the epidural space can be difficult using the conventional loss of resistance technique. Acoustic amplification has been used to detect puncture more accurately. An in-line pressure transducer is attached to a pressure amplifier, which in turn is attached to a loudspeaker. The sound of the needle passing from low resistance surrounding tissues to the high resistance of the ligamentum flavum can be appreciated audibly (Lechner et al, 2002).

## **Complications and contraindications**

In general safe practice, knowing when a local anesthetic technique is contraindicated is essential, as is an understanding of the complications of local anesthetic administration and how to treat them.

Complications can be broadly categorized as local or systemic problems.

## Local

### *Skin*

Local analgesic techniques should not be attempted where the skin overlying the site of injection is inflamed, infected, or clearly neoplastic as this may promote transmission and spread of infectious organisms or neoplastic cells into adjacent tissue.

### *Nerve*

Direct trauma to nerves from the needle or high-pressure injection may result in loss of sensation, pain, discomfort and/or motor weakness that may be temporary, lasting days to years, or permanent. Damage to blood vessels close to nerves, particularly those with poor blood supply, may lead to ischemia. Finally, local anesthetic drugs or additives may be directly toxic to nervous tissue.

## Systemic

### *Central nervous system*

Excessive systemic uptake of local anesthetic can cause seizures. This occurs because inhibitory neurons are more sensitive to  $\text{Na}^+$  channel blockade, and if they are blocked then excitation results.

### *Cardiovascular*

Local anesthetic techniques should be performed with caution or avoided in patients that are hypotensive or in shock, as inadvertent drug administration into a vein or unexpected rapid systemic drug absorption may lead to cardiovascular collapse.

### *Coagulopathy*

Techniques in which hemorrhage is a possible complication, e.g. epidural block, should not be performed in patients where coagulopathy has been demonstrated or is suspected. Laceration of a blood vessel may lead to bleeding that will be hard to control. If uncontrolled hemorrhage occurs in a closed space, it may lead to pressure necrosis of adjacent tissues.

## Species differences and considerations

As a species, cats are generally less amenable to being restrained for procedures unless they are heavily sedated or anesthetized, compared to dogs in which some techniques can be performed with light sedation. It can also be more challenging to accurately weigh cats, which may lead to inadvertent under- or overdosage. Regardless of species,



accurate dosing will lead to fewer complications. Inhalant anesthesia with isoflurane or sevoflurane decreases metabolism of lidocaine after 2 mg/kg intravenous administration in cats, even at relatively low inhalant doses, whereas this is not the case in dogs (Thomasy et al, 2005). Rapid inadvertent intravenous injection of lidocaine, while not ideal in any species, may be more problematic in cats.

## Safe practice

### Gaining confidence

Local anesthetic techniques vary in their level of difficulty to perform, effectiveness of block, and complexity of equipment required. Starting by using straightforward techniques that require little in the way of extra equipment, and have been reported to have high success rates, is recommended. Once comfortable with simple techniques, the practitioner can expand his or her repertoire. Attending continuing education seminars and courses on local anesthetic techniques, particularly when a wet lab is a component of the training, is recommended. We also believe that using this book will be an asset when it comes to performing techniques where the practitioner has little or no experience.

### Making a plan

Regardless of technique, successfully performing local anesthetic blocks requires preparation of the appropriate equipment, and a thorough understanding of patient anatomy and local anesthetic drug pharmacology. It is also important that the clinician performing the local anesthetic block has an appreciation of the potential complications that might occur and is able to develop a plan to treat them (see Chapter 7). The person performing the block should also have in mind an alternative plan for providing analgesia should the block prove impossible to perform, or in cases where continuing with the block is contraindicated (e.g. severe drop in blood pressure during epidural administration, skin infection over injection site becomes apparent after clipping).

## References

- Chu, C.R., Izzo, N.J., Coyle, C.H., et al. (2008) The in vitro effects of bupivacaine on articular chondrocytes. *J Bone Joint Surg* **90**: 814–820.
- Lechner, T.J.M., van Wijk, M.G.F. & Maas, J.J. (2002) Clinical results with a new acoustic device to identify the epidural space. *Anaesthesia* **57**: 768–772.

- Riley, E.T. & Carvalho, B. (2007) The Episure syringe: a novel loss of resistance syringe for locating the epidural space. *Anesth Analg* **105**: 1164–1166.
- Thomasy, S.M., Pypendop, B.H., Ilkiw, J.E. & Stanley, S.D. (2005) Pharmacokinetics of lidocaine and its active metabolite, monoethylglycinexylidide, after intravenous administration of lidocaine to awake and isoflurane-anesthetized cats. *Am J Vet Res* **66**: 1162–1166.

## Further reading

- Campoy, L. & Read, M.R. (2013) *Small Animal Regional Anesthesia and Analgesia*. Ames, IA: Wiley-Blackwell.
- Shelby, A.M. & McKune, C.M. (2014) *Small Animal Anesthesia Techniques*. Ames, IA: Wiley-Blackwell.
- Tranquilli, W.J., Thurmon, J.C. & Grimm, K.A. (2007) *Lumb & Jones Veterinary Anesthesia*, 4th edn. Ames, IA: Wiley-Blackwell.

## CHAPTER 2

# Cutaneous Innervation Index

Gwen Covey-Crump


---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.  
By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.  
© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## Cutaneous innervation index


Use this chapter as an aid to choosing an appropriate regional anesthetic technique for the planned procedure.

### Canine dermatomes – Body




**Figure 2.1** Canine dermatome map. Cutaneous innervation according to spinal segment. This is intended as a guide only as much overlap between segments and variation between animals exist. Redrawn from Oliver, J.E. & Lorenz, M.D. (eds) (1983) *Veterinary Neurologic Diagnosis*. Philadelphia: W.B.Saunders Co. Reproduced with permission from The Ohio State University.

## Head



**Figure 2.2** Cutaneous innervation of the canine head. Autonomous sensory regions are shown. Some overlap and variation between zones and animals exist. Reproduced with permission from The Ohio State University.

Thoracic limb




**Figure 2.3** Cutaneous innervation of the canine thoracic limb. Autonomous sensory regions are shown. Some overlap and variation between zones and animals exist. LCB-T<sub>2</sub>, lateral cutaneous brachial nerve-thoracic 2. Reproduced with permission from The Ohio State University.

**Table 2.1** Sensory innervation of the canine thoracic limb

	Region	Nerve	Techniques
C6–7	Lateral shoulder and joint	Suprascapular	CP
		Subscapular	CP
C7–8	Craniomedial antebrachium	Musculocutaneous	CP, BP, RUMM
C8	Caudal shoulder joint, lateral brachium	Axillary	CP, BP
C7–T1	Lateral elbow joint, craniolateral antebrachium, carpus and digits	Radial	CP, BP, RUMM
C8–T1	Medial elbow joint	Median	CP, BP, RUMM
C8–T2	Caudal elbow joint, caudolateral antebrachium, carpus and digits	Ulnar	CP, BP, RUMM

BP, brachial plexus; CP, cervical paravertebral; RUMM, radial, ulnar, median, musculocutaneous.

Pelvic limb



**Figure 2.4** Cutaneous innervation of the canine pelvic limb. Autonomous sensory regions are shown. Some overlap and variation between zones and animals exist. Reproduced with permission from The Ohio State University.



**Table 2.2** Sensory innervation of the canine pelvic limb

Region				Techniques	
L4	Hemipelvis (excl. skin overlying hip joint), medial thigh, stifle, leg, hock	Femoral	Saphenous	Inguinal Preiliac Psoas compartment	
L5					
L6					
L7	Entire limb distal to stifle (excl. medial strip)	Dorsolateral distal limb	Sciatic	Peroneal	Lateral Parasacral
S1, S2		Laterocaudal distal limb		Tibial	Transgluteal

## CHAPTER 3

# Infiltration Blocks

Fernando Martinez Taboada

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.  
By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.  
© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## **Infiltration for mass removal**

### **Line and ring blocks**

#### **Indications**

Surgical resection of cutaneous or superficial masses, surgical closure of lacerations, procedures involving an appendage (e.g. toe, tail, pinna, etc.)

#### **Target nerves**

Cutaneous nerves of the affected area

#### **Region anesthetized**

Appendage distal to the block, area immediately underneath the block and, in the case of performing a ring block, area circumscribed by it

#### **Landmarks**

Surrounding tissue around the area of interest. It is fundamental that the subcutaneous tissue is clearly identified

#### **Needle**

Hypodermic needle. It is recommended to use a longer length but small in size needle (e.g. 23 Ga 2.5–3.8 cm)

#### **Depth**


Subcutaneous injection of local anesthetic

#### **Technique**


- 1 Identify the area of interest. If a mass is being resected, the total contour of it and the margins must be taken into account
- 2 The line and ring blocks are the easiest blocks to perform, as they do not require specific knowledge of neuroanatomy
- 3 Introduce a long hypodermic needle to its whole length in the subcutaneous tissue
- 4 Aspirate to prevent accidental intravenous administration and deposit some local anesthetic (e.g. 0.2 mL bupivacaine)
- 5 Withdraw the needle approximately 5 mm (the distance can vary depending on the volume injected in each point. The bigger the volume, the longer the distance can be), aspirate again and inject a bit more local anesthetic
- 6 Repeat the sequence until the area of interest is circumscribed
- 7 When performing ring or curve line blocks, it is recommended to withdraw the needle almost totally before redirecting it in a different direction to prevent tissue damage and needle bending

#### **Cautions**

Accidental intravenous injection of local anesthetic. This type of block substitutes for accuracy by using a large volume of local anesthetic. Overdose of local anesthetic is a serious hazard if the total dose is not considered. With this in mind, if the volume is insufficient, use of a more dilute preparation is recommended



**Figure 3.1** Line block. Reproduced with permission from The Ohio State University.



**Figure 3.2** Ring block. Performed as a combination of line blocks to avoid needle bending. Reproduced with permission from The Ohio State University.

## **Inverted “L” block**

### ***Indications***

Surgical resection of cutaneous or superficial masses, surgical closure of lacerations

### ***Target nerves***

Cutaneous nerves originating in the dorsal spinal nerves

### ***Region anesthetized***

The area of interest usually is located within the trunk of the animal. Classically, this technique has been used to desensitize the flank

### ***Landmarks***

Surrounding tissue around the area of interest. It is fundamental to identify the subcutaneous tissue and the muscular layers, if there is an involvement of these structures

### ***Needle***

Hypodermic needle. It is recommended to use a longer length but small in size needle (e.g. 23 Ga 2.5–3.8 cm)

### ***Depth***

Subcutaneous injection of local anesthetic

### ***Technique***

- 1 Identify the area of interest. If a mass is being resected, the total contour of it and the margins must be taken into account
- 2 Following the steps described in the previous block, two line blocks must be performed (forming the inverted “L”)
- 3 Perform a line block between the mass (or area of interest) and the patient’s spine
- 4 Add another line block cranial to the area of interest
- 5 If there is involvement of muscular layers, it is necessary to block the subcutaneous tissue (for blocking the cutaneous branches of the nerves), but also to perform a deep block to block the muscular branches. The injection of local anesthetic in the muscle is very stimulating, so the subcutaneous block is advisable before attempting the deeper block

### ***Cautions***

Accidental intravenous injection of local anesthetic. Overdose of local anesthetic is a serious hazard as a great volume of local anesthetic may be necessary to desensitize a large area of the body. Dilution of the local anesthetic with sterile saline may be an option to achieve the desired volume



**Figure 3.3** Inverted “L” block (*blue arrows*) for the resection of mast cell tumor (*red arrow*).

## Block of the surgical incision

This block can be achieved by a single injection at the time of closure or via a diffusion catheter placed in the deep layers of the surgical incision allowing continuous administration of local anesthetics.

### Single injection of the surgical incision

#### *Indications*

Analgesia of wounds or surgical incisions

#### *Target nerves*

Cutaneous nerves of the affected area

#### *Region anesthetized*

Surgical incision edges

#### *Landmarks*

Subcutaneous tissue around the incision

#### *Needle*

23–25 Ga 1.6–1.9 cm hypodermic needle

#### *Depth*

Skin to subcutaneous tissue


#### *Technique*

- 1 Probably the oldest and easiest nerve block to perform
- 2 The injection of local anesthetic in the subcutaneous tissue exposed in the surgical wound is more effective than a simple splash of the drug in the incision
- 3 Injection of local anesthetic should be done in the same way as the line block (see previous description in this chapter)

Savvas et al (2008) reported better results when the block was performed before the first incision than when it was performed before suturing.

#### *Cautions*

Overdose of local anesthetic is a serious hazard as a great volume of local anesthetic may be necessary to desensitize a large area of the body. Dilution of the local anesthetic with sterile saline may be an option to achieve the desired volume



**Figure 3.4** Single injection incisional block. Reproduced with permission from The Ohio State University.



## **Diffusion catheter/"soaker" catheter/"pain buster"**

### **Indications**

Analgesia of wounds or surgical incisions

### **Target nerves**

Cutaneous nerves of the affected area

### **Region anesthetized**

Surgical incision from deep layers to skin edges

### **Landmarks**

Surgical incision

### **Needle**

None. The catheter is a flexible polyurethane tube of small diameter with multiple pores for even distribution of the local anesthetic. The catheter is placed during the surgical closure of the incision

### **Depth**

Skin to the deepest layers of the surgical wound


### **Technique**

- 1 Using a pair of artery forceps and a scalpel, introduce the catheter a few centimeters from the incision edge (so it does not interfere with healing)
- 2 Place the catheter in the deeper layers of the incision
- 3 Make sure the length of tube with pores is completely in the incision and that it is long enough for the size of the incision
- 4 Close the soft tissue above the catheter
- 5 Secure the catheter externally to the skin to prevent its movement
- 6 The analgesia can be achieved by the intermittent administration of bupivacaine 2 mg/kg every 6 h (some overlapping between administrations is required as local anesthetics can sting during administration). A constant infusion avoids an analgesia "gap" between doses. Constant infusion dose of lidocaine: dogs 2–3 mg/kg/h, cats 1–1.5 mg/kg/h

Hansen et al (2013) demonstrated no difference between commercial catheters and handmade ones. They also found that very slow infusion rates produce an erratic distribution of the drug.

### **Cautions**

The total volume to be used is dependent on the incision length and not the size of the patient. Always double-check the total dose to be injected and dilute it with sterile saline, if the volume is not sufficient.



**Figure 3.5** Placement of a soaking catheter in the deep layers of the incision before closure. Reproduced with permission from The Ohio State University.



**Figure 3.6** Soaking catheter used in a cat for pain management after an injection site sarcoma.

## Intercostal blocks

### *Indications*

Lateral thoracotomy, analgesia for rib fractures, analgesia for chest drain placement

### *Target nerves*

Intercostal nerves

### *Region anesthetized*

Intercostal space distal to the injection

### *Landmarks*

Caudal edge of the rib as proximal to the spine as is possible to palpate

### *Needle*

23 Ga 1.9–2.5 cm

### *Depth*

Caudal edge of the rib. *Caution:* introducing the needle further may penetrate the pleura. Directing the tip of the needle underneath the rib may damage the intercostal vein and/or artery


### *Technique*

- 1 Palpate the caudal edge of the rib and follow it towards the spine
- 2 Introduce the needle perpendicular to the skin and direct its tip towards the caudal edge of the rib
- 3 Aspirate to prevent intravascular injection
- 4 Administer 0.3–1 mL bupivacaine
- 5 Repeat the previous steps in at least two intercostal spaces cranial and caudal to the area of interest (Thompson & Johnson, 1991). Although desensitizing three spaces, the results are better


This technique has been shown to be superior to systemic opioids when respiratory function was assessed in patients after lateral thoracotomy (Berg & Orton, 1986).

### *Cautions*

Hematoma, hemothorax, intravascular injection, nerve damage



**Figure 3.7** Intercostal nerve block. Palpate the rib and then slide the needle over the caudal edge. Reproduced with permission from The Ohio State University.



**Figure 3.8** An effective intercostal block requires three intercostal spaces on either side of the area of interest. Reproduced with permission from The Ohio State University.

## Interpleural block (also known as intrapleural or pleural)

### Indications

Classically, this block is indicated for the postoperative management of a lateral thoracotomy or a sternotomy, as it produces less respiratory depression than opioids alone (Berj & Orton, 1986; Conzemius et al, 1994). In human medicine, it is indicated for other conditions, e.g. brachial plexus tumor (Dionne, 1992), cholecystectomy (VadeBoncouer et al, 1989), but veterinary medicine lacks experience in these uses.

### Target nerves

The mechanism of action is unknown. It is speculated that the technique may work at three levels: intercostal nerves (due to a retrograde diffusion through the parietal pleura), blockage of the sympathetic chain and the splanchnic nerves, and/or blockage of the ipsilateral brachial plexus.

### Region anesthetized

Thoracic cavity, cranial abdominal cavity

### Landmarks

Interpleural cavity in an intercostal space

### Needle

Over-the-needle catheter (14–18 Ga 4.8–3.2 cm). It is advisable to use catheters with bayonet tips to prevent puncture of the lung

### Depth

Skin to interpleural cavity


### Technique

For repeatable block, it is advisable to place an indwelling thoracic drain or diffusion catheter although the technique may be performed by single injection.

- 1 Introduce the catheter through the skin over an intercostal space and advance the tip of the needle, tunneling in the subcutaneous tissue to the following intercostal space
- 2 Penetrate the thoracic wall at the intercostal space, avoiding the caudal edge of the rib where the nerve, artery, and vein are located
- 3 Introduce the catheter at a shallow angle to decrease the chance of damaging the lung
- 4 Analgesia can be provided with bupivacaine 2 mg/kg every 5–6 h

### Cautions

Pneumothorax, hemothorax, overdose of local anesthetic. Complications are rare if the catheter/chest drain is manipulated carefully. The cardiopulmonary effects of the local anesthetics in the interpleural space are very transient and not clinically significant, even after pericardectomy (Bernard et al, 2006)



**Figure 3.9** Interpleural block. The needle is introduced closer to the cranial edge of the rib to avoid the intercostal vessels. Reproduced with permission from The Ohio State University.



**Figure 3.10** Over-the-needle catheter with bayonet tip to prevent lung damage.

## Intraperitoneal block

### *Indications*

Analgesia of the abdominal cavity. This block has become a common postoperative analgesic technique for laparotomies and laparoscopies in human medicine

### *Target nerves*

Splanchnic nerves responsible for the visceral component of the abdominal pain

### *Region anesthetized*

Abdominal cavity (exclusively the visceral component)

### *Landmarks*

Abdominal cavity (specifically the area of interest, e.g. ovary, kidney, gallbladder ...)

### *Needle*

No needle is required, it is applied in the abdomen before closing the linea alba

### *Depth*

Abdominal cavity

### *Technique*

- 1 Use 1–2 mg/kg bupivacaine
- 2 Dilute with sterile saline if the area of interest is too large for the calculated volume
- 3 Splash the local anesthetic over the area of interest, for instance the ovarian pedicle
- 4 Alternatively, splash the volume into the cavity, trying to achieve a uniform distribution
- 5 Close linea alba

Some authors recommend infiltration of the surgical incision to minimize parietal or somatic pain (Carpenter et al, 2004).

### *Cautions*

The total volume to be used is dependent on the abdominal cavity or the size of the area of interest and not so much on the length and size of the patient. Always double-check the total dose to be injected and dilute it with sterile saline, if the volume is not sufficient.

## References

- Berg, R.J. & Orton, E.C. (1986) Pulmonary function in dogs after intercostal thoracotomy: comparison of morphine, oxymorphone, and selective intercostal nerve block. *Am J Vet Res* **47**: 471–474.
- Bernard, F., Kudnig, S.T. & Monnet, E. (2006) Hemodynamic effects of interpleural lidocaine and bupivacaine combination in anesthetized dogs with and without open pericardium. *Vet Surg* **35**: 252–258.
- Carpenter, R.E., Wilson, D.V. & Evans, A.T. (2004) Evaluation of intraperitoneal and incisional lidocaine or bupivacaine for analgesia following ovari-hysterectomy in the dog. *Vet Anaesth Analg* **31**: 46–52.
- Conzemius, M.G., Brockman, D.J., King, L.G., et al. (1994) Analgesia in dogs after intercostals thoracotomy: clinical trial comparing intravenous buprenorphine and interpleural bupivacaine. *Vet Surg* **23**: 291–298.
- Dionne, C. (1992) Tumour invasion of the brachial plexus: management of pain with interplural analgesia. *Can J Anaesth* **39**: 520–521.
- Hansen, B., Lascelles, B.D., Thomson, A., et al. (2013) Variability of performance of wound infusion catheters. *Vet Anaesth Analg* **40**(3): 308–315.
- Savvas, I., Papazoglou, L.G., Kazakos, G., et al. (2008) Incisional block with bupivacaine for analgesia after celiotomy in dogs. *J Am Anim Hosp Assoc* **44**(2): 60–66.
- Thompson, S.E. & Johnson, J.M. (1991) Analgesia in dogs after intercostal thoracotomy. A comparison of morphine, selective intercostals nerve block, and interpleural regional analgesia with bupivacaine. *Vet Surg* **20**: 73–77.
- VadeBoncouer, T.R., Riegler, F.X., Gautt, R.S., et al. (1989) A randomized, double blind comparison of the effects of interpleural bupivacaine and saline on morphine requirements and pulmonary function after cholecystectomy. *Anesthesiology* **71**: 339–343.



## CHAPTER 4

# Blocks of the Head

Fernando Martinez Taboada

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.

By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.

© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## Retrobulbar

### *Indications*

Surgical procedures involving the eye and/or procedures requiring akinesia of the eye

### *Target nerves*

Ophthalmic division of the trigeminal nerve (V) (sensory perception), also within the ocular cone, cranial nerves optic (II), oculomotor (III) and abducens (VI) (motor function of the extraocular muscles). The trochlear nerve (IV) is outside the ocular cone, but is frequently blocked by diffusion of the anesthetic drugs and contributes to the akinesia of the eye

### *Region anesthetized*

Eye including conjunctiva, cornea, and uvea

### *Landmarks*

Eye globe, orbit of the eye, zygomatic arch, vertical ramus of the mandible

### *Needle*

22 Ga 2.5 cm (up to 3.8 cm, depending on animal size)

### *Depth*

Skin to ocular cone

### *Technique*

- 1 Locate the depression limited by the caudal aspect of the maxilla, the ventral edge of the zygomatic arch and the vertical ramus of the mandible
- 2 Introduce the needle perpendicular to the long axis of the head
- 3 As soon as the needle tip is in the subcutaneous tissue, direct the needle mediodorsal (behind the zygomatic arch) towards the ocular cone
- 4 Accurate location of the tip of the needle can be confirmed as the pressure of the needle against the extraocular muscles produce a ventrodorsal movement of the globe
- 5 Aspirate to prevent intravascular injection
- 6 Inject 1–2 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)

### *Cautions*

Hematoma, intravascular injection, nerve damage. Intraocular damage is rare, although intraconal hematomas can cause blindness. The traditional approach for this block, curving a spinal needle (20 Ga 7.5 cm), has been associated with failure of the technique and risk of direct subarachnoid injection. For this reason, this alternative approach is presented here, although further research comparing differences in safety is still needed



**Figure 4.1** (a) Insertion of the needle in the ocular cone produces ventrodorsal movement of the globe. (b–c) Retrobulbar block (subzygomatic approach). Parts (a) and (c) reproduced with permission from The Ohio State University.

## Maxillary

### *Indications*

Surgical procedures involving the maxilla, upper lip and/or palate, dental extractions (Aguiar et al, 2015) and rhinoscopy (if performed bilaterally) (Cremer et al, 2013)

### *Target nerves*

Maxillary nerve (ideally with its branches: the nasal and major palatine nerves)

### *Region anesthetized*

Dorsal head and muzzle approximately up to midline. Maxilla and maxillary teeth, hard and soft palate (if major palatine nerve is blocked with the technique), nares, upper labium and rostral part of the nasal cavity

### *Landmarks*

Zygomatic arch, caudal part of the maxilla, and vertical ramus of the mandible

### *Needle*

22 Ga 2.5 cm non-insulated needle

### *Depth*


Skin to pterygoid fossa

### *Technique*

- 1 Identify the depression formed by the caudal aspect of the maxilla, the ventral edge of the zygomatic arch, and vertical ramus of the mandible. This depression is located ventral to the lateral canthus of the eye
- 2 Introduce the needle perpendicular to the maxilla
- 3 Direct the tip of the needle perpendicular to the sagittal plane of the skull
- 4 If the tip of the needle hits bone, withdraw approximately 1–2 mm before performing the injection
- 5 Aspirate to prevent intravascular injection
- 6 Inject 0.3–1 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine or ropivacaine)

### *Cautions*

Hematoma, intravascular injection, nerve damage



**Figure 4.2** (a) Maxillary nerve block. Needle is directed perpendicular to the pterygoid fossa to also block the nasal and palatal nerve branches. (b) Maxillary nerve block. Part (a) reproduced with permission from The Ohio State University.

## Infraorbital

### *Indications*

Surgical procedures involving the rostral part of the muzzle (some procedures may require bilateral block)

### *Target nerves*

Infraorbital nerve

### *Region anesthetized*

Rostral part of the muzzle including skin, maxilla, maxillary teeth, and palate. Rostral upper lip and nasal cavity

### *Landmarks*

Infraorbital foramen, premolar tooth P3

### *Needle*

25 Ga 1.5 or 2.5 cm hypodermic needle

### *Depth*


Skin to the interior of the infraorbital foramen

### *Technique*

- 1 Percutaneously or lifting the upper lip, locate the infraorbital foramen (dorsal to the tooth P3)
- 2 Introduce the needle at a shallow angle towards the entrance of the foramen
- 3 Advance the needle into the foramen as far as possible (within a safety margin, a few millimetres is sufficient)
- 4 Aspirate to avoid intravascular injection
- 5 Inject 0.3–0.5 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)
- 6 The effectiveness of the block and the size of the area desensitized depend on whether the injection is performed inside the foramen (anesthetizing up to the first molar; Gross et al, 1997) or only at the entrance (blocking only the area rostral to the canine teeth)

### *Cautions*

Hematoma, intravascular injection, nerve damage



**Figure 4.3** (a) Nerve branches exiting the infraorbital foramen. (b) Infraorbital block (note the insertion of the needle tip within the canal). (c) Infraorbital block. Parts (a) and (b) reproduced with permission from The Ohio State University.

## **Mandibular (or inferior alveolar)**

### **Indications**

Surgical procedures involving the mandible, lower lip, and dental extractions (Aguilar et al, 2015)

### **Target nerves**

Inferior alveolar nerve (at the mandibular foramen)

### **Region anesthetized**

Hemi-mandible up to the chin. The tissues between the mandibles should not be anesthetized if the technique is performed correctly. On some occasions, the most rostral part of the chin is not desensitized (in some animals the innervation is by the mylohyoid nerve)

### **Landmarks**

Mandible, angular process, mandibular foramen

### **Needle**

22 Ga 1.6 or 2.5 cm

### **Depth**

Skin rostral to the angular process of the mandible to the mandibular foramen


### **Technique**

- 1 Palpate the angular process of the mandible, then open the mouth and palpate the mandibular foramen. This foramen is located approximately halfway between the last molar tooth and the condylar process
- 2 Introduce the needle percutaneously immediately rostral to the angular process and directed dorsally along the medial aspect of the mandible
- 3 Direct the tip of the needle towards the foramen. It is very difficult to advance the needle into the foramen, but the block can be effective as long as the anesthetic solution is delivered at the entrance of the foramen
- 4 Aspirate to prevent intravascular injection
- 5 Inject 0.2–1 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)

### **Cautions**

Hematoma, intravascular injection, nerve damage. Desensitization of the tongue can occur if a bilateral block is performed with large volumes and/or poor technique (difficulty locating and injecting within the foramen or as close as possible to it). This desensitization may lead to self-mutilation of the tongue





**Figure 4.4** (a) Innervation of the mandible and the tongue. (b) Percutaneous approach to the mandibular nerve (note the needle tip close to the foramen). (c) Percutaneous approach for the mandibular nerve block. Parts (a) and (b) reproduced with permission from The Ohio State University.

## Mental (or middle mental)

### *Indications*

Surgical procedures involving the rostral mandible, teeth associated with the area, and rostral lower lip

### *Target nerves*

Inferior alveolar (at the mental foramen; also known as the mental nerve)

### *Region anesthetized*

Chin when performed bilaterally, up to premolar tooth P4 if injection is performed inside the foramen (Gross et al, 2000)

### *Landmarks*

Mental foramen, lower lip, premolar tooth P2

### *Needle*

25 Ga 1.6 mm

### *Depth*


Gum mucosa to foramen

### *Technique*

- 1 Palpate the foramen ventral to the tooth P2 from the inside of the lower lip
- 2 Introduce the needle just rostral to that point, at a shallow angle, towards the foramen
- 3 Advance the needle into the foramen as far as possible (without forcing it in)
- 4 Aspirate to prevent intravascular injection
- 5 Inject 0.2–0.6 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)

### *Cautions*

Hematoma, intravascular injection, nerve damage. In some instances, the most rostral part of the chin is not desensitized (in some animals the innervation is by the mylohyoid nerve)



**Figure 4.5** (a) Mental block (note the insertion of the needle tip within the nerve canal). (b) Mental block for rostral hemimandibulectomy. Part (a) reproduced with permission from The Ohio State University.

## **Auriculotemporal and great auricular**

### ***Indications***

Analgesia in severe cases of otitis externa, perioperative analgesia for surgical interventions affecting the external ear canal and the auricular pinna (postoperative analgesia can also be achieved by local infiltration at the time of closure, but the use of this block would provide intraoperative analgesia as well)

### ***Target nerves***

Auriculotemporal (branch of the mandibular nerve) and great auricular (formed by the union of the ventral roots of C1 and C2)

### ***Region anesthetized***

External ear canal and pinna

### ***Landmarks***


Zygomatic arch, external ear canal, vertebral atlas transverse process (also known as wings)

### ***Needle***

22 Ga 2.5 cm (or longer in big dogs)

### ***Depth***

Skin to mid-distance between skin and temporomandibular joint (auriculotemporal nerve) and skin to subcutaneous tissue (great auricular nerve)



**Figure 4.6** (a) Innervation of the ear and pinna. (b) Identification of the anatomical landmarks for the desensitization of the auriculotemporal nerve.

## **Technique**

### **Auriculotemporal nerve block**


- 1** Palpate the external ear canal (it is conical in shape and very hard when inflamed)
- 2** Locate the caudal aspect of the zygomatic arch
- 3** Introduce the needle perpendicular to the skin in the palpable depression between the zygomatic arch and the external ear canal
- 4** Advance the needle as far as possible until the tip is touching bone (the temporomandibular joint)
- 5** Knowing the total depth of the tissues, withdraw the needle approximately halfway
- 6** Aspirate to prevent intravascular injection
- 7** Inject 0.5–1.5 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)

### **Great auricular nerve**

- 1** Palpate the external ear canal
- 2** Locate the most cranial point of the transverse process of the atlas vertebra
- 3** Introduce the needle parallel to the transverse process
- 4** Direct the tip of the needle ventral to the most cranial part of the process
- 5** The nerve is very superficial and it is better to maintain the needle at a very shallow angle to the skin
- 6** Aspirate to prevent intravascular injection
- 7** Inject 0.5–1.5 mL of anesthetic solution or combination (e.g. 0.25–0.5% bupivacaine)

## **Cautions**

Hematoma, intravascular injection, nerve damage. Temporary paralysis of the facial and/or temporopalpebral nerves preventing the animal from blinking (recommended to lubricate the eye every 2–4 h)



(c)



(d)

**Figure 4.6** (continued)

(c) Innervation in relation to anatomical landmarks. (d) Great auricular nerve block. Parts (a) and (c) reproduced with permission from The Ohio State University.

## References

- Aguiar, J., Chebroux, A., Martinez-Taboada, F., et al. (2015) Analgesic effects of maxillary and inferior alveolar nerve blocks in cats undergoing dental extractions. *J Feline Med Surg* **17**: 110–116.
- Cremer, J., Sum, S.O., Braun, C., et al. (2013) Assessment of maxillary and infraorbital nerve blockade for rhinoscopy in sevoflurane anesthetized dogs. *Vet Anaesth Analg* **40**: 432–439.
- Gross, M.E., Pope, E.R., O'Brien, D., et al. (1997) Regional anesthesia of the infraorbital and inferior alveolar nerves during noninvasive tooth pulp stimulation in halothane-anesthetized dogs. *J Am Vet Med Assoc* **211**: 1403–1405.
- Gross, M.E., Pope, E.R., Jarboe, J.M., et al. (2000) Regional anesthesia of the infraorbital and inferior alveolar nerves during noninvasive tooth pulp stimulation in halothane-anesthetized cats. *Am J Vet Res* **61**: 1245–1247.



## CHAPTER 5

# Regional Anesthetic Blocks of the Limbs

Gwen Covey-Crump

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.

By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.

© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## **Thoracic limb regional anesthetic blocks**

**Table 5.1** Elicited motor responses to nerve stimulation – thoracic limb

Technique	Motor response	Muscle	Nerves	Accept	Comment
Cervical paravertebral	Hiccup	Diaphragm	Phrenic	✗	Too ventral/caudal
	Adduct, rotate scapula/shoulder	Supraspinatus, infraspinatus	Suprascapular	✗/	Cranial
			Subscapular		
brachial plexus	Rotate shoulder in, flex elbow	Biceps brachialis	Musculocutaneous	✓	Cranial aspect of technique
	Flex shoulder, extend elbow/carpus	Teres major/minor, deltoideus	Axillary	✓	Middle aspect of technique
	Extend elbow/carpus	Triceps, carpal/digital extensors	Radial	✓/	Local anesthetic here should diffuse over most of brachial plexus
	Pronate antebrachium, carpal/palmar flexion	Carpal/digital flexors	Median	✓	Injection very caudal may not reach more cranial nerves
	Palmar flexion	Carpal/digital flexors	Ulnar	✓	
RUMM	Palmar extension	Carpal/digital extensors	Radial	✓	
	Palmar flexion	Carpal/digital flexors	Median, ulnar	✓	
	Triceps/biceps contraction			✗	Direct muscle stimulation

## Brachial plexus

### *Indications*

Unilateral surgical procedures of distal humerus, elbow, antebrachium, carpus

### *Target nerves*

Musculocutaneous, axillary, radial, median and ulnar

### *Region anesthetized*

Thoracic limb distal to mid humerus

### *Landmarks*

Acromion, trachea, jugular vein, 1<sup>st</sup> rib

### *Needle*

21 Ga 5.0–10.0 cm insulated needle

### *Depth*


Acromion to 1<sup>st</sup> rib

### *Technique (Campoy et al, 2008; Mahler & Adogwa, 2008)*

- 1 Identify 1<sup>st</sup> rib by axillary palpation. Premeasure needle depth to a point where a line extending the course of the jugular vein crosses the 1<sup>st</sup> rib. The brachial plexus is just cranial to this point
- 2 Insert needle craniomedial to the acromion in a caudoventral direction parallel to the jugular vein and in a strictly sagittal plane
- 3 With nerve stimulator at 2 mA and gradually reducing, observe responses to nerve stimulation (Table 5.1) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 4 At point of radial nerve stimulation (and no further than premeasured depth), aspirate to avoid intravascular (blood) or interpleural (air) injection
- 5 Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine
- 6 Withdraw needle 0.5–1 cm, aspirate, then inject further 0.1 mL/kg
- 7 Repeat step 6, injecting a final 0.1 mL/kg beneath the point of the shoulder

### *Cautions*

Intravascular, interpleural injection, hematoma. Nerve damage



**Figure 5.1** Brachial plexus block. (a) Anatomy, (b) approach. Part (a) reproduced with permission from The Ohio State University.

## Paravertebral: C6–T1

### Indications

Unilateral surgical procedures of the scapula, shoulder, and brachium, including forequarter amputation

### Target nerves

Spinal nerves of C6, C7, C8, T1

### Region anesthetized

Entire thoracic limb, except skin over upper shoulder region

### Landmarks

Transverse process C6, head of rib 1

### Needle

21 Ga 5.0 cm insulated (very small dogs) to 15.0 cm (very large dogs)


### Technique (Campoy & Read, 2013; Lemke & Creighton, 2008)

Advanced technique to be undertaken with due caution only in patients where palpation of C6 and 1<sup>st</sup> rib is possible. Unilateral block only

- 1 With patient in lateral recumbency and limb to be blocked uppermost, retract scapula caudally
- 2 Palpate transverse process of C6, maintain digital pressure on lateral aspect
- 3 From dorsolateral approach, advance needle in caudal direction to meet transverse process. Walk needle off cranial border
- 4 With nerve stimulator at 2 mA and gradually reducing, observe responses to nerve stimulation of ventral branches of spinal nerves C6 and C7, suprascapular, subscapular, musculocutaneous (Table 5.1) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 5 Aspirate to avoid intravascular (blood) or spinal (clear fluid) injection
- 6 Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine
- 7 Repeat steps 3–6, walking needle off caudal border of transverse process
- 8 Palpate axillary artery at 1<sup>st</sup> rib costochondral junction and thoracic inlet. Advance needle to meet cranial aspect 1<sup>st</sup> rib 1–2 cm dorsal to costochondral junction
- 9 Observe elbow/carpus/digit extension (C8) and carpus/digit flexion (T1)
- 10 Aspirate to avoid intravascular (blood) or interpleural (air) injection
- 11 Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine

### Cautions

Intravascular epidural or interpleural injection, hematoma. Nerve injury. Phrenic nerve paralysis (common). Do not perform technique bilaterally



**Figure 5.2** Paravertebral block. (a) Anatomy, (b) approach. Part (a) reproduced with permission from The Ohio State University.

## **RUMM (radial/ulnar/median/musculocutaneous)**

### **Indications**

Anesthesia for surgical procedures of the thoracic limb distal to and including the carpus

### **Target nerves**

Radial (R), ulnar (U), median (Me), musculocutaneous (Mus) nerves

### **Region anesthetized**

Carpus, metacarpus, and digits

### **Landmarks**

Lateral and medial epicondyles, greater tubercle of humerus. Lateral and medial head of triceps, biceps and biceps brachialis muscles. Brachial artery

### **Needle**

21 Ga 5.0 cm insulated/20–22 Ga 2.0–4.0 cm non-insulated

### **Depth**

10–20 mm


### **Technique (Lamont & Lemke, 2008; Trumpatori et al, 2009)**

- 1 Begin with patient in lateral recumbency, limb to treat uppermost, elbow flexed 90°
- 2 Grasp humerus from cranial aspect, place thumb on caudal humerus two-thirds distance from greater tubercle to lateral epicondyle. Retract brachialis muscle cranially. Insert needle from caudal aspect, perpendicular to long axis of humerus at 45° angle. Puncture long heads of triceps to contact humerus, withdraw slightly
- 3 If using nerve stimulator, with nerve stimulator at 2 mA and gradually reducing, observe responses to radial nerve stimulation (Table 5.1) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 4 Aspirate to avoid intra-arterial injection
- 5 Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine
- 6 Turn patient into opposite lateral recumbency, have assistant retract upper limb caudally (limb to treat is lowest and flexed 90°)
- 7 Palpate brachial artery at point mid humerus. With digital pressure on humerus, retract biceps brachialis muscle and brachial artery cranially. Insert needle from caudal aspect perpendicular to long axis of humerus and at 45° angle to contact humerus, withdraw slightly
- 8 Observe responses to median/ulnar nerve stimulation if using (Table 5.1)
- 9 Aspirate to avoid intra-arterial injection
- 10 Inject half of 0.15 mL/kg 0.5% bupivacaine or 0.75% ropivacaine deeply and remaining half as needle is withdrawn

### **Cautions**

Risk of intravascular injection, nerve damage





**Figure 5.3** Radius, ulnar, median, musculocutaneous (RUMM) block. (a) Anatomy, (b) approach. Part (a) reproduced with permission from The Ohio State University.

## **Bier block (intravenous regional anesthesia) (Staffieri, 2013)**

### ***Indications***

Surgical procedures of <90 min duration of the distal limb, especially when a bloodless surgical field is advantageous

### ***Target nerves***

Local anesthetic effect by spread to peripheral nerve endings

### ***Region anesthetized***

Thoracic limb distal to elbow and pelvic limb distal to tarsus

### ***Important anatomy***

- Suitable distal vein
- Palpable peripheral artery

### ***Equipment***


- Tourniquet – preferably pneumatic with pressure gauge
- Esmarch or elastic bandage
- Doppler probe (optional)
- Catheter 22–25 Ga 25–33 mm

### ***Technique (Webb et al, 1999)***

- 1 Clip surgical and catheter placement sites (plus tourniquet site if heavily coated)
- 2 Place intravenous catheter as distally as possible
- 3 Find a palpable artery and mark with pen
- 4 Apply tourniquet and identify lower occlusion pressure (LOP) (pressure at which palpable pulse is absent)
- 5 Release tourniquet, apply esmarch or elastic bandage tightly
- 6 Apply tourniquet to 50–100 mmHg above LOP, record time
- 7 Release esmarch bandage
- 8 Confirm absence of peripheral pulse
- 9 Infuse lidocaine 0.5% 0.6 mL/kg over 2–3 min, observe patient for toxicity
- 10 Remove catheter
- 11 During surgery, monitor and maintain cuff pressure 50–100 mmHg above systolic arterial pressure
- 12 At end of surgery, remove tourniquet slowly, observing for signs of toxicity and to ensure hemostasis

### ***Cautions***

Avoid cuff pressures in excess of 400 mmHg. Use only lidocaine. Risk of systemic toxicity, ischemic injury, limb engorgement, tourniquet pain



**Figure 5.4** Bier block. (a) Find lower occlusion pressure (LOP), (b) apply esmarch bandage, (c) apply tourniquet 50–100 mmHg above LOP and remove esmarch, (d) check pulse absent, infuse lidocaine.

## **Pelvic limb**

**Table 5.2** Elicited motor responses to nerve stimulation – pelvic limb

Technique	Motor response	Muscle	Nerves	Accept	Comment
Femoral – inguinal approach	Patellar twitch	Quadriceps	Femoral/saphenous	✓	
	Anterior thigh	Sartorius		✗	Too superficial
Femoral – lateral preiliac approach (caudal lumbar plexus)	Hip flexion	Psoas stimulation		✗	Direct muscle stimulation
	Hip adduction	Gracilis, pectineus, adductor femoris, obturator externus	Obturator	✗	May block with femoral
	Stifle extension, patellar twitch	Quadriceps	Femoral	✓	
	Biceps twitch at 1 mA			✗	Too superficial
Sciatic – lateral approach	Semimembranosus/tendinosus twitch		Direct muscle stimulation	✗	Too superficial
	Stifle flexion	Biceps femoris	Motor branches only	✗	Too caudal
	Plantarflexion	Gastrocnemius	Sciatic	✓	
	Dorsiflexion		Tibial	✓	
			Peroneal/fibular	✓	
	Gluteal twitch			✗	Direct muscle stimulation
Sciatic – parasacral approach	Stifle flexion	Biceps femoris	Sciatic	✓	
	Plantarflexion	Gastrocnemius	Tibial	✓	
	Dorsiflexion		Peroneal/fibular	✓	
				✓	

## **Femoral/saphenous nerve block**

### ***Inguinal approach***

#### ***Indications***

Surgery of distal femur, stifle and distal limb in combination with sciatic nerve block

#### ***Target nerves***

Femoral and/or its branch: saphenous

#### ***Region anesthetized***

Medial aspect of hindlimb – mid femur to 1st digit

#### ***Landmarks***

Femoral artery; femoral triangle (enclosed dorsally by rectus femoris, cranially by caudal belly of sartorius muscle, proximally (deep) by iliopsoas muscle)

#### ***Needle***

21 Ga 5.0 cm insulated/20–22 Ga 2.0–4.0 cm non-insulated

#### ***Depth***


5–15 mm

#### ***Technique***

- 1** With patient in lateral recumbency, hindlimb to treat uppermost
- 2** Abduct limb caudally, palpate femoral artery
- 3** Direct needle dorsomedially towards nerve which lies cranial to femoral artery
- 4** With nerve stimulator at 2 mA and gradually reducing, observe responses to nerve stimulation including quadriceps twitch, patellar twitch, stifle extension (Table 5.2) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection) (sartorius twitch = needle too superficial)
- 5** Aspirate to avoid intra-arterial injection
- 6** Inject 0.2 mL/kg 0.5% bupivacaine or 0.75% ropivacaine

#### ***Cautions***

Hematoma, intravascular injection, nerve damage



**Figure 5.5** Femoral nerve block – inguinal approach. (a) Anatomy, (b) approach. Part (a) reproduced with permission from The Ohio State University.

## ***Lateral preiliac approach (caudal lumbar plexus)*** ***(Portela et al, 2013)***

### ***Target nerves***

Femoral (may also block obturator)

### ***Region anesthetized***

Hemipelvis, femur, medial femorotibial joint, skin of dorsomedial tarsus, 1st digit. May not anesthetize skin overlying coxofemoral joint

### ***Landmarks***

Cranial dorsal iliac crest (CDIC), L6

### ***Needle***

Small dogs 22 Ga 5.0 cm, large dogs 22 Ga 7.5 cm insulated

### ***Depth***


Use depth of iliopsoas muscle as reference

### ***Technique***

- 1 With patient in lateral recumbency, hindlimb to treat uppermost
- 2 Draw a line dorsoventral from L6, bisect with line parallel to spine originating from most cranial aspect of iliac crest
- 3 Puncture site at bisection, advance needle caudomedially with 30–45° intent through iliocostalis lumborum muscle
- 4 With nerve stimulator at 1 mA and gradually reducing, observe responses to nerve stimulation – quadriceps contraction (Table 5.2) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 5 Aspirate to avoid intra-arterial injection
- 6 Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine



### ***Cautions***

Low risk of epidural migration, vessel puncture



**Figure 5.6** (a) Puncture site.





**Figure 5.6** Femoral nerve block – lateral preiliac approach. (b) Anatomy, (c) approach. Part (a) reproduced with permission from The Ohio State University.

## Sciatic nerve block

### *Indications*

Surgery of distal femur, stifle and distal limb in combination with femoral nerve block

### *Target nerves*

Sciatic (ischiatric) nerve gives off branches: common peroneal, tibial nerves

### *Region anesthetized*

Lateral, dorsal, caudal aspects of hindlimb – caudolateral stifle to digits 2–5

### *Lateral approach (Campoy et al, 2008)*

### *Landmarks*

Greater trochanter of femur (GT), ischial tuberosity (IT)

### *Needle*

21 Ga 5.0 cm insulated or 22 Ga 2.0–4.0 cm non-insulated

### *Depth*


5–10 mm

### *Technique*

- 1 With patient in lateral recumbency, hindlimb to treat uppermost on a line between GT and IT, puncture site is 1/3 distance from GT
- 2 Direct needle perpendicular to skin
- 3 With nerve stimulator at 2 mA and gradually reducing, observe responses to nerve stimulation – dorsiflexion of tarsus (peroneal nerve), plantarflexion (tibial nerve) (Table 5.2) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 4 Aspirate to avoid intra-arterial injection
- 5 Inject 0.05–0.1 mL/kg 0.25–0.5% bupivacaine or 0.75% ropivacaine

### *Cautions*

Risk of intravascular injection, nerve damage



**Figure 5.7** Sciatic nerve block – lateral approach. (a) Anatomy, (b) approach. Part (a) reproduced with permission from The Ohio State University.

## ***Parasacral approach (sacral plexus block) (Portela et al, 2010)***

### ***Landmarks***

Cranial dorsal iliac crest (CDIC), ischial tuberosity (IT), dorsal spinous processes (DSP)

### ***Needle***

21 Ga 5.0 cm insulated

### ***Depth***


15–30 mm

### ***Technique***

- 1** With patient in lateral recumbency, hindlimb to treat uppermost, on a line between CDIC and IT, puncture site is 1/3 distance from CDIC
- 2** Direct needle sagittally
- 3** With nerve stimulator at 2 mA and gradually reducing, observe responses to nerve stimulation – gastrocnemius twitch, digital/tarsal flexion/extension (Table 5.2) (reducing to 0.2 mA to avoid intrafascicular injection prior to injection)
- 4** Aspirate to avoid intra-arterial injection
- 5** Inject 0.1 mL/kg 0.5% bupivacaine or 0.75% ropivacaine

### ***Cautions***

Risk of intravascular injection, epidural injection, nerve damage, bowel puncture



**Figure 5.8** Sciatic nerve block – parasacral approach. (a) Anatomy, (b) approach. Reproduced with permission from The Ohio State University.

## References

- Campoy, L. & Read, M.R. (2013) The thoracic limb. In: *Small Animal Regional Anesthesia and Analgesia*. Ames, IA: Wiley-Blackwell, pp.146–147.
- Campoy, L. Looney, A.L. Ludders, J.W. & Gleed, R.D. (2008) Distribution of a Lidocaine-methylene blue solution staining in brachial plexus, lumbar plexus and sciatic nerve blocks in the dog. *Vet Anaesth Analg* **35**: 348–354.
- Lamont, L.A. & Lemke, K.A. (2008) The effects of medetomidine on radial nerve blockade with mepivacaine in dogs. *Vet Anaesth Analg* **35**: 62–68.
- Lemke, K.A. & Creighton, C.M. (2008) Paravertebral blockade of the brachial plexus in dogs. *Vet Clin Small Anim* **38**: 1231–1241
- Mahler, S.P. & Adogwa, A.O. (2008) Anatomical and experimental studies of brachial plexus, sciatic, and femoral nerve-location using peripheral nerve stimulation in the dog. *Vet Anaesth Analg* **35**: 80–89.
- Portela, D.A., Otero, P.E. Tarragona, L., Briganti, A., Breggi, G. & Melanie, P. (2010) Combined paravertebral plexus block and parasacral sciatic block in healthy dogs. *Vet Anaesth Analg* **37**: 531–541.
- Portela, D.A., Otero, P.E., Briganti, A., et al. (2013) Femoral nerve block: a novel psoas compartment lateral pre-iliac approach in dogs. *Vet Anaesth Analg* **40**: 194–204.
- Staffieri, F. (2013) Intravenous regional anesthesia. In: Campoy, L. & Read, M.R. (eds) *Small Animal Regional Anesthesia and Analgesia*. Ames, IA: Wiley-Blackwell, pp. 261–271.
- Trumpatori, B.J., Carter, J.E., Hash, J., et al. (2009) Evaluation of a mid-humeral block of the radial, ulnar, musculocutaneous and median (RUMM block) nerves for analgesia of the distal aspect of the thoracic limb in dogs. *Vet Surg* **39**: 785–796.
- Webb, A.A., Cantwell, S.L., Duke, T., et al. (1999) Intravenous regional anesthesia (Bier block) in a dog. *Can Vet J* **40**: 419–421.

## CHAPTER 6

# Epidurals and Spinals

Turi K. Aarnes

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.  
By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.  
© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## **Lumbosacral epidural and spinal in dogs**

### **Indications**

Abdominal pain, thoracic pain, hindlimb pain, perineum/tail

### **Target nerves**

Nerves caudal to the spread of local anesthetic

### **Region of anesthesia/analgesia**

Using 0.2 mL/kg of injectate (local anesthetic or morphine at appropriate dose diluted to appropriate volume), anesthesia/analgesia should occur from T13 spinal space and extend caudally. Larger volumes may be used if cranial spread is desired.

### **Landmarks**

Place the thumb and middle finger on the craniodorsal aspects of the wings of the ilium and place the index finger at the lumbosacral space (L7–S1) for epidural administration (Figure 6.1). When performing a spinal, place the thumb and middle finger on the craniodorsal aspects of the wings of the ilium, and the index finger at the space between L5 and L6.

### **Needle**

A spinal needle with stylette (e.g. Tuohy needle for epidural, Quincke needle for spinal), depending on the size of the dog – typically a 20 Ga, 5.1 cm needle for a 15–40 kg dog.

### **Technique (Skarda & Tranquilli, 2007a; Troncy et al, 2002)**

A distinct “pop” is felt when the needle is advanced through the interarcuate ligament (ligamentum flavum) (Figure 6.2). No blood or spinal fluid should be observed when performing an epidural. Fluid will most likely be observed flowing from the needle when the stylette is removed when performing a spinal, though it is possible to be in the subarachnoid space with no fluid flowing (“dry tap”). Loss of resistance technique may be utilized by injecting approximately 1 mL air – no resistance to injection and no backpressure on the syringe plunger should be felt. The “hanging drop technique” may also be used by placing saline into the needle to fill the hub. The drop should be “sucked in” to the epidural space. Minimal or no resistance to injection should be felt during drug administration.


### **Advantages**

Advantages of epidural anesthesia include good muscle relaxation, postoperative analgesia, minimal effects on the body, low cost





**Figure 6.1** Palpation of the lumbosacral space in a dog. Place the thumb and middle finger on the craniodorsal aspects of the wings of the ilium and place the index finger at the lumbosacral space.



**Figure 6.2** Lumbosacral epidural technique in a dog. A distinct “pop” is felt when the needle is advanced through the interarcuate ligament (ligamentum flavum). Reproduced with permission from The Ohio State University.

### **Complications**

Inadequate anesthesia due to faulty technique, if the animal is awake during surgery (might move); hypotension (if using local anesthetic), respiratory depression (local anesthetic epidurals and spinals), or apnea after excessive blockade (epidurals and spinals); accidental administration into the subarachnoid space (if intent is to perform an epidural).

### **Contraindications**

Hypovolemia and hypotension (when using local anesthetics), bleeding disorder, skin infection

## **Lumbosacral epidural in cats**

### **Indications**

Abdominal pain, thoracic pain, hindlimb pain, perineum/tail

### **Target nerves**

Nerves caudal to the spread of local anesthetic

### **Region of anesthesia/analgesia**

Using 0.2 mL/kg of injectate (local anesthetic or morphine at appropriate dose diluted to appropriate volume), anesthesia/analgesia should occur from T13 spinal space and extend caudally. Local anesthetic for lumbosacral epidural should be used with caution in cats due to toxicity/sensitivity and potential for spinal administration.

### **Landmarks**

Place the thumb and middle finger on the craniodorsal aspects of the wings of the ilium and place the index finger at the lumbosacral space (Figure 6.3).

### **Needle**

A 22 Ga 2.5 cm straight needle

### **Technique (Skarda & Tranquilli, 2007b; Troncy et al, 2002)**

A 22 Ga needle is placed at the midline of the lumbosacral space (L7–S1) (Figure 6.4). A distinct “pop” is felt when the needle is advanced through the interarcuate ligament (ligamentum flavum). No blood or spinal fluid should be observed. Minimal resistance to injection should be felt.

### **Complications**


Inadequate anesthesia due to faulty technique, if the animal is awake during surgery (might move); hypotension, respiratory depression, or apnea after excessive blockade; accidental administration into the subarachnoid space. Due to anatomical differences from dogs, more likely to perform a spinal rather than an epidural (spinal cord ends at L6–S2), in which case dose of injected drug should be reduced by 50%.

### **Contraindications**

Hypovolemia and hypotension (when using local anesthetics), bleeding disorder, skin infection. Reduce dose by 50% if performing spinal rather than epidural.



**Figure 6.3** Palpation of the lumbosacral space in a cat. Place the thumb and middle finger on the craniodorsal aspects of the wings of the ilium and place the index finger at the lumbosacral space.



**Figure 6.4** Lumbosacral epidural and spinal technique in a cat. A 22 Ga needle is placed at the midline of the lumbosacral (L7–S1) space. Reproduced with permission from The Ohio State University.

## Caudal epidural in dogs and cats

### *Indications*

Described for use in cats with urethral obstruction. Other possibilities: dystocia, tail amputation, perineal procedures

### *Target nerves*

Nerves caudal to the spread of local anesthetic

### *Region of anesthesia/analgesia*

Perineum, tail, sacrum

### *Landmarks*

A coccygeal epidural can be performed at either the sacrococcygeal interspace or Co1–Co2. First moveable vertebrae between the sacrum and tail, or first coccygeal interspace, on midline

### *Needle*

25 Ga 1.5 cm straight needle

### *Technique (O'Hearn & Wright, 2011)*


Palpate the space between sacrum and Co1 or Co1 and Co2. Insert 25 Ga needle on midline, direct at a 30–45° angle, and advance through the ligamentum flavum (Figure 6.5). Aspirate through the needle to check for blood (if blood is aspirated, start over). Infuse 0.1–0.2 mL/kg of lidocaine 2%.

### *Complications*

Bleeding, incomplete blockade

### *Contraindications*

Hypovolemia and hypotension (when using local anesthetics), bleeding disorder, skin infection



**Figure 6.5** Coccygeal epidural technique in a cat. Insert a 25 Ga needle on the midline, directed at a 30–45° angle, and advance through the ligamentum flavum. Reproduced with permission from The Ohio State University.

## References

- O'Hearn, A.K. & Wright, B.D. (2011) Coccygeal epidural with local anesthetic for catheterization and pain management in the treatment of feline urethral obstruction. *J Vet Emerg Crit Care* **21**: 50–52.
- Skarda, R.T. & Tranquilli, W.J. (2007a) Local and regional anesthetic and analgesic techniques: dogs. In: Tranquilli, W.J., Thurmon, J.C. & Grimm, K.A. (eds) *Lumb & Jones' Veterinary Anesthesia and Analgesia*, 4th edn. Ames, IA: Wiley-Blackwell, pp.561–594.
- Skarda, R.T. & Tranquilli, W.J. (2007b) Local and regional anesthetic and analgesic techniques: cats. In: Tranquilli, W.J., Thurmon, J.C. & Grimm, K.A. (eds) *Lumb & Jones' Veterinary Anesthesia and Analgesia*, 4th edn. Ames, IA: Wiley-Blackwell, pp.595–603.
- Troncy, E., Junot, S., Keroack, S., et al. (2002) Results of preemptive epidural administration of morphine with or without bupivacaine in dogs and cats undergoing surgery: 265 cases (1997–1999). *J Am Vet Med Assoc* **221**: 666–672.

## CHAPTER 7

# Emergency Procedures

Turi K. Aarnes

---

*Handbook of Small Animal Regional Anesthesia and Analgesia Techniques*, First Edition.  
By Phillip Lerche, Turi K. Aarnes, Gwen Covey-Crump and Fernando Martinez Taboada.  
© 2016 John Wiley & Sons, Ltd. Published 2016 by John Wiley & Sons, Ltd.

## **Complications associated with local anesthetic techniques**

### **Bleeding**

Bleeding is a potential complication of all nerve blocks. Caution should be exercised in patients with bleeding/clotting disorders and patients that have been heparinized. Epidurals and spinals are contraindicated in these patients.

### *Clinical signs*

Visible hemorrhage, hypotension

### *Treatment*

Monitor systemic blood pressure, apply pressure to area of hemorrhage, cold compress to reduce blood flow, in extreme cases ligation of blood vessels (e.g. accidental laceration of an artery)

### **Nerve trauma**

Injection of local anesthetic directly into nerve or nerve trauma may result from local anesthetic techniques. Return to normal function may occur over time.

### *Clinical signs*

Pain, motor or sensory deficits, urinary or fecal incontinence

### *Treatment*

Supportive care, systemic analgesics for pain, systemic anti-inflammatories

## **Complications associated with local anesthetic drugs**

### **Allergic reactions to local anesthetic drugs**

True allergic reactions to local anesthetics are rare. Reported with use of aminoesters (e.g. procaine, tetracaine). Signs develop 12–24 h after exposure (Lin & Liu, 2013).

### *Clinical signs*

Hypersensitivity reaction; contact dermatitis including erythema, pruritus, papules, vesicles



### **Treatment**

Stop administration of local anesthetic, airway maintenance, epinephrine, intravenous fluids, diphenhydramine

### **Methemoglobinemia**

More likely with benzocaine or prilocaine but also reported with procaine and lidocaine. Also more likely in cats than dogs.

### **Clinical signs**

Hypoxia, cyanosis, death, blood that looks brown in color, dyspnea, nausea, tachycardia

### **Treatment**

Intravenous administration of 1% methylene blue; intubation and oxygen supplementation, ventilation, ECG monitoring (Stoelting & Hillier, 2006).

### **Systemic toxicity of local anesthetics**

Systemic toxicity of local anesthetics is usually associated with increased plasma concentrations. Systemic toxicities may result in cardiac arrest. Monitoring patients and preparation for performing CPR are necessary.

### **Cardiovascular toxicity** (Table 7.1)

Cardiovascular (CV) toxicity is increased with bupivacaine compared with lidocaine or ropivacaine, accidental intravenous administration of bupivacaine or ropivacaine, and in patients with decreased hepatic function. Large doses of high-potency local anesthetics are more likely to result in CV toxicity leading to complete AV block and CV collapse. Separately, CV toxicity may result from CNS toxicity due to decreased activity of the nucleus tractus solitarius resulting in decreased autonomic control of the CV system (Salinas et al, 2004).

### **CNS toxicity** (Table 7.2)

Related to plasma concentration and potency. Low concentrations cause CNS depression, high concentrations can result in excitation due to inhibition of inhibitory neurons. High potency and high lipid solubility can result in increased CNS toxicity (Lin & Liu, 2013).

Table 7.1 Cardiovascular toxicity

Complication	Mechanism	Clinical presentation/clinical sign	Treatment
Vasodilation	Excessive plasma concentration, also related to potency of local anesthetic drug	Hypotension, collapse, tachycardia	Stop administration of local anesthetic, continuous monitoring of invasive blood pressure, administer IV fluids, vasopressors (vasopressin, epinephrine, ephedrine, phenylephrine), inotropes (dopamine, dobutamine), intravenous lipid emulsion
Dysrhythmias	Excessive plasma concentration, unbound drug diffuses to conducting tissue of heart, also related to potency of local anesthetic drug	ECG changes: prolongation of P-R interval, prolongation of QRS interval, AV block, ventricular premature contractions, ventricular tachycardia, ventricular fibrillation	Stop administration of local anesthetic, continuous monitoring of ECG, intravenous lipid emulsion, antidysrhythmics, oxygen supplementation
Myocardial depression and cardiac toxicity	Unbound drug diffuses to conducting tissue of heart, inhibition of depolarization rate of cardiac action potential due to sodium channel blockade; most likely with accidental IV administration of bupivacaine  Epinephrine and phenylephrine may increase bupivacaine toxicity to cAMP inhibition	ECG changes, hypotension	Stop administration of local anesthetic, continuous monitoring of invasive blood pressure, continuous ECG monitoring, intravenous fluids, administer IV fluids, vasopressors (vasopressin, epinephrine, ephedrine, phenylephrine), inotropes (dopamine, dobutamine), intravenous lipid emulsion (Weinberg et al, 2003), antidysrhythmics, oxygen supplementation

Table 7.2 CNS toxicity

Complication	Mechanism	Clinical presentation/clinical sign	Treatment
CNS depression/toxicity	Excessive plasma concentration (dose dependent), also related to potency of local anesthetic drug	Nystagmus, myoclonus, seizures	Stop administration of local anesthetic, anticonvulsants: midazolam or diazepam 0.5 mg/kg IV; intravenous lipid emulsion; oxygen administration
	Excessive plasma concentration (dose dependent), also related to potency of local anesthetic drug	Unconsciousness, coma	Stop administration of local anesthetic, continuous monitoring of ECG, monitor ventilation, airway maintenance, oxygen supplementation, epinephrine, intravenous fluids, intravenous lipid emulsion
	Excessive plasma concentration (dose dependent), also related to potency of local anesthetic drug	Respiratory arrest	Stop administration of local anesthetic, intubation and oxygen supplementation, ventilation, intravenous lipid emulsion
Neurotoxicity	Rare complication. Direct exposure of nerves to local anesthetics can result in toxicity of neural fibers, leading to concentration- dependent nerve damage and decreased neural blood flow. More likely to see neurotoxicity with spinals and epidurals. Return to normal function may occur over time	Pain, motor or sensory deficits, urinary or fecal incontinence	Stop administration of local anesthetic, supportive care, systemic analgesics for pain

## References

- Lin, Y. & Liu, S.S. (2013) Local anesthetics. In: Barasch, P.G., Cullen, B.F., Soteling, R.K., et al. (eds) *Clinical Anesthesia*, 7th edn. Philadelphia: Lippincott Williams & Wilkins, pp.561–577.
- Salinas, F.V., Liu, S.L. & Scholz, A.M. (2004) Analgesics: ion channel ligands/sodium channel blockers/local anesthetics. In: Evers, A.S. & Maze, M. (eds) *Anesthetic Pharmacology: Physiologic Principles and Clinical Practice*. Philadelphia: Churchill Livingstone, pp.507–533.
- Stoelting, R.K. & Hillier, S.C. (2006) Local anesthetics. In: Stoelting, R.K. & Hillier, S.C. (eds) *Pharmacology & Physiology in Anesthetic Practice*, 4th edn. Philadelphia: Lippincott Williams & Wilkins, pp.179–207.
- Weinberg, G., Ripper, R., Feinstein, D.L., et al. (2003) Lipid emulsion infusion rescues dogs from bupivacaine-induced cardiac toxicity. *Reg Anesth Pain Med* **28**: 198–202.

## Further reading

- Martin-Flores, M. (2013) Clinical pharmacology and toxicology of local anesthetic and adjuncts. In: Campoy, L. & Read, M.R. (eds) *Small Animal Regional Anesthesia and Analgesia*. Ames, IA: Wiley-Blackwell, pp.25–38.
- Skarda, R.T. & Tranquilli, W.J. (2007) Local anesthetics. In: Tranquilli, W.J., Thurmon, J.C. & Grimm, K.A. (eds) *Lumb & Jones' Veterinary Anesthesia and Analgesia*, 4th edn. Ames, IA: Wiley-Blackwell, pp.395–418.

# Index

- acromion, 56
- additives, 6, 10
- adrenaline, 6
- allergic reactions, 84–85
- alpha-2 adrenoceptor
  - agonists, 2, 6
- amino amide, 3
- amino ester, 3, 84
- amputation, 58
- antebrachium, 55–56
- auriculotemporal nerve
  - block, 48–50
  - ear, 48–50
  - ear canal, 48, 50
  - otitis, 48
  - pinna, 48–49
  - zygomatic arch, 38–40, 48, 50
- Bier block, 6, 62–63
- block
  - auriculotemporal, 48–50
  - Bier block, 62–63
  - brachial plexus, 56–57
  - diffusion, 28
  - femoral/saphenous, 66–69
  - great auricular, 48–50
  - incisional, 26
  - infraorbital, 42–43
  - intercostal, 30–32
  - interpleural, 32–33
  - intraperitoneal, 34
  - inverted “L”, 24–25
  - line, 22–24
  - mandibular, 44–45
  - maxillary, 40–41
  - mental, 46–47
  - paravertebral C6–T1, 58–59
  - radial, ulnar, median,
    - musculocutaneous (RUMM), 17, 60–61
  - retrobulbar, 38–39
  - ring block, 22–24
  - sacral plexus block, 72–73
  - sciatic block, 70–71
- blood vessels
  - axillary a., 59
  - brachial a., 60–61
  - brachial v., 61
  - caudal gluteal a. v., 71
  - cephalic v., 61
  - external iliac a. v., 68
  - femoral a. v., 66–67
  - internal iliac a. v., 68
  - jugular v., 56, 59
- brachial plexus, 17, 55–57
- bupivacaine, 3, 5–7
- carpus, 55–56, 58, 60
- caudal epidural, 80
- coxofemoral joint, 68
- cranial dorsal iliac crest, 72–73
- dexmedetomidine, 6
- diaphragm, 55
- diffusion catheter, 28–29
  - catheter, 28–29
  - pain buster, 29
  - soaker, 28–29
  - soaker catheter, 28–29
- digit, 6, 58, 60, 66, 68, 70

diphenhydramine, 85  
dorsal spinous process, 72

elbow, 55–56, 58, 60, 62  
epidural, 7  
epidural catheter, 8  
epinephrine, 5–6, 85  
esmark bandage, 62–63

femoral triangle, 66  
femorotibial joint, 68–69  
forequarter, 58

great auricular nerve  
    block, 48, 50  
    ear, 48–50  
    ear canal, 48, 50  
    otitis, 48  
    pinna, 48–49  
greater trochanter, 70–71

humerus, 56, 60  
hyaluronidase, 7

iliac crest, 68, 72  
infiltration of surgical incision,  
    22–34

infraorbital nerve block, 42–43  
    dental, 40–44  
    muzzle, 40, 42  
    nares, 40

intercostal block, 30–33  
    chest wall, 30–33  
    intercostal, 30–33  
    rib, 30–33

interpleural block, 32–33  
    chest drain, 32–33

intraperitoneal  
    abdominal cavity, 34  
    laparoscopy, 34  
    laparotomy, 34

inverted “L” block, 25–28  
ischial tuberosity, 70–72

ketamine, 2, 6

laceration, 22–28  
lidocaine, 3–6, 11, 80  
line block, 22–23, 27–28  
lumbosacral epidural, 76

mandibular nerve block, 44–45  
    dental, 44  
    mandibulectomy, 44–47  
mass removal, 22, 24, 26  
maxillary nerve block, 40–41  
    dental, 40  
    pterygoid fossa, 40  
    rhinoscopy, 40  
    zygomatic arch, 40

mechanism of action, 4  
medetomidine, 6  
mental nerve block, 46–47  
    chin, 46–47  
    dental, 46–47

mepivacaine, 3, 5  
metabolism, 4, 11  
metacarpus, 60  
methemoglobinemia, 85  
methylene blue, 85  
morphine, 6

muscle  
    adductor femoris m., 65, 67  
    biceps brachii m., 55, 60–61  
    biceps femoris m., 65  
    brachialis m., 60–61  
    carpal extensors, 55  
    carpal flexors, 55  
    deltoideus m., 55  
    digital extensors, 55  
    digital flexors, 55  
    gastrocnemius m., 65, 72  
    gracilis m., 65, 67  
    iliopsoas m., 66–69  
    obturator externus m., 65  
    pectineus m., 65, 67  
    psoas m., 65  
    quadriceps m., 65–66, 68  
    rectus femoris m., 66–67  
    sartorius m., 65–67  
    semimembranosus m., 65

- semitendinosus m., 65
- teres major m., 55
- teres minor m., 55
- triceps, 55, 60
- N-methyl-D-aspartate (NMDA)
  - channel, 6
- needles
  - hypodermic, 7
  - Quincke, 76
  - spinal, 7
  - Tuohy, 7–8, 76
- nerve
  - auriculotemporal, 48–50
  - axillary n., 17, 55–57
  - brachiocephalic n., 15
  - cervical n., 6–8, 17, 57
  - common peroneal n., 70
  - cutaneous
    - C2, 14
    - C3, 14
    - LCB-T<sub>2</sub>, 15
  - facial n., 14
  - femoral n., 19, 65–69
  - great auricular, 48–50
  - infraorbital n., 42–43
  - infraspinatus n., 55
  - ischiatric n., 70
  - lumbar plexus, 65, 68
  - mandibular n., 14, 44–45
  - maxillary n., 14, 40–41
  - median n., 17, 55–57, 60–61
  - musculocutaneous n., 15, 17, 55–57, 60–61
  - obturator n., 65, 68
  - ophthalmic n., 14, 38
  - peroneal n., 16, 19, 65, 70
  - phrenic n., 55, 58
  - radial n., 15, 17, 55–57, 60–61
  - sacral plexus, 72–73
  - saphenous n., 16, 19, 65–66
  - sciatic n., 16, 19, 65, 70–73
  - spinal nerves, 58
  - subscapular n., 17, 55, 58
  - suprascapular n., 17, 55, 58
  - supraspinatus n., 55
  - thoracic n., 1, 57, 59
  - tibial n., 16, 19, 65, 70
  - ulnar n., 15, 17, 55–56, 60–61
- nerve locator
  - acoustic device, 9
  - peripheral nerve stimulator, 8
  - pressure transducer, 9
  - ultrasound, 9
- nociception, 2, 6
- opioids, 2, 6
- patellar twitch, 65–66
- pelvic limb
  - femoral/saphenous nerve
    - block, 65–69
  - inguinal approach, 65–67
  - lateral pre-iliac approach
    - (caudal lumbar plexus), 65, 68–69
  - parasacral approach (sacral plexus block), 65, 72–73
  - sciatic nerve block, 65–66, 70–73
- pK<sub>a</sub>, 4–5
- preemptive analgesia, 2
- prilocaine, 3
- proparacaine, 3
- retrobulbar block, 38–39
  - eye, 38–39
  - ocular, 38–39
  - zygomatic arch, 38–40, 48, 50
- rib, 58
- ring block, 6
- ropivacaine, 3, 5, 7
- route of administration
  - epidural, 4, 6–11, 76–81
  - intra-articular, 4
  - intravenous regional, 3, 5
  - peripheral nerve block, 4
  - regional infiltration, 3
  - spinal, 2, 4, 6–8, 76–79

- route of administration (*Continued*)
  - topical
    - corneal, 3
    - intrapleural, 3
    - transdermal, 3
- scapula, 55, 58
- shoulder, 55–56, 58
- sodium bicarbonate, 7
- spinal, 76–77
- spine, 76–81
- stifle, 65–66, 70
- syringes, 3, 7–8
- tarsus, 62, 68–70
- tourniquet, 62–63
- toxicity
  - cardiovascular, 85
  - CNS (central nervous system), 85
- transverse process, 58
- xylazine, 6
- zygomatic arch, 38–40, 48, 50